arXiv:1704.04806v1 [math.ST] 16 Apr 2017

Simultaneous Inference for High Dimensional Mean Vectors

By Zhipeng Lou and Wei Biao Wu

Department of Statistics, University of Chicago

April 18, 2017

Abstract

Let $X_1, \ldots, X_n \in \mathbb{R}^p$ be i.i.d. random vectors. We aim to perform simultaneous inference for the mean vector $\mathbb{E}(X_i)$ with finite polynomial moments and an ultra high dimension. Our approach is based on the truncated sample mean vector. A Gaussian approximation result is derived for the latter under the very mild finite polynomial ($(2+\theta)$-th) moment condition and the dimension p can be allowed to grow exponentially with the sample size n. Based on this result, we propose an innovative resampling method to construct simultaneous confidence intervals for mean vectors.

1 Introduction

Let $X_1, \ldots, X_n$ be i.i.d. random vectors in $\mathbb{R}^p$ with mean vector $\mathbb{E}X_i = \mu$ and covariance matrix $\mathrm{Cov}(X_i) = \Sigma$. We are interested in conducting statistical inference for μ when the dimension p can be comparable to or even much larger than n. Estimating μ by the traditional sample mean $\hat{\mu} = \sum_{i=1}^n X_i/n$, Chernozhukov et al. (2013) and Chernozhukov et al. (2014) stated that under suitable moment conditions, as $n \to \infty$ and possibly $p = p_n \to \infty$,

$$\rho_n := \sup_{t \in \mathbb{R}} \left| \mathbb{P}(\sqrt{n}|\hat{\mu} - \mu|_\infty \le t) - \mathbb{P}(|Y|_\infty \le t) \right| \to 0. \tag{1}$$

where $Y = (Y_1, \ldots, Y_p)^\top \in \mathbb{R}^p$ is a centered Gaussian vector with $\mathrm{Cov}(Y) = \Sigma$ and $|x|_\infty = \max_{j \le p} |x_j|$ is the usual ℓ_∞ norm for a vector $x = (x_1, \ldots, x_p)^\top \in \mathbb{R}^p$. Suppose we only have uniformly finite q-th ($q > 3$) moment for each coordinate of X_i, i.e., there exists a constant $C > 0$ such that

$$\max_{j \le p} \mathbb{E}|X_{ij}|^q \le C \tag{2}$$

for some $q > 3$. We can show that the condition

$$p(\log p)^{3q/2-1} = o(n^{q/2-1}) \tag{3}$$

is nearly optimal for (1); see Proposition 2.2. If (3) is barely violated, then under (2) we can have $\rho_n \to 1$; see (12) and (13). Hence in general, the allowed dimension p can be at most a power of n if we use $\hat{\mu}$ for inference of μ. In this paper, we propose a new approach to perform simultaneous inference for the mean vector with finite polynomial moments and show that our method applies under ultra high dimensional settings, in which $\log p = o(n^c)$ for some $c > 0$.

In Section 2, we shall study properties of the truncated sample mean with element-wise truncation. A similar idea was adopted by Fan et al. (2016) in the estimation of covariance matrices. We shall establish a Gaussian approximation result for the truncated sample mean vector with uniformly finite $(2+\theta)$-th $(0 < \theta \leq 1)$ moments for X_{ij}, $1 \leq j \leq p$. As an important feature, the dimension p can be as large as $e^{o(n^c)}$ for some $c > 0$.

Equipped with the Gaussian approximation result for the truncated sample mean vector, we are ready to perform simultaneous statistical inference for μ. In particular, we propose an innovative resampling method called truncated half sampling procedure to construct simultaneous confidence intervals of μ in Section 3. As a main advantage of this method, we do not need to deal with the problem of estimating the covariance matrices which is highly nontrivial and computationally intensive in the high dimensional case, and which may require extra structural assumptions.

2 The truncated sample mean vector and a Gaussian approximation theory

Given any $\kappa > 0$, let $t_\kappa : \mathbb{R} \to [-\kappa, \kappa]$ be the truncated function defined by $t_\kappa(x) = (x \wedge \kappa) \vee (-\kappa)$. Given the data $X_1, X_2, \ldots, X_n$, define the truncated sample mean vector

$$\hat{\mu}_\kappa = \frac{1}{n} \sum_{i=1}^{n} t_\kappa(X_i), \tag{4}$$

where $t_\kappa(X_i) = (t_k(X_{i1}), \ldots, t_\kappa(X_{ip}))^\top \in \mathbb{R}^p$. With properly chosen truncated level κ, we establish a Gaussian approximation result for $|\hat{\mu}_\kappa|_\infty$ with uniformly finite $(2+\theta)$-th

moments for X_{ij}, $1 \le j \le p$.

Theorem 2.1. *Let $\mu = \mathbb{E}X_i = 0$. Assume there exist constants $b, M_\theta > 0$ such that $\min_{j\le p} \mathbb{E}|X_{ij}|^2 \ge b$ and $M_\theta = \max_{j\le p} \mathbb{E}|X_{ij}|^{2+\theta} < \infty$ for some $0 < \theta \le 1$. Further assume*

$$(\log p)^{4+3\theta} M_\theta^2 = o(n^\theta). \tag{5}$$

and we take $\kappa \asymp (nM_\theta/\log p)^{1/(2+\theta)}$. Then as $n \to \infty$,

$$\rho_{n,\kappa} := \sup_{t\in\mathbb{R}} |\mathbb{P}(\sqrt{n}|\hat{\mu}_\kappa|_\infty \le t) - \mathbb{P}(|Y|_\infty \le t)| \to 0. \tag{6}$$

Proof. For simplicity of notation, write $\mathbb{E}_0[X] = X - \mathbb{E}X$. Elementary calculation implies for $q = 1, 2$,

$$\max_{j\le p} \mathbb{E}|X_{ij}|^q \mathbf{1}\{|X_{ij}| \ge \kappa\} \le \kappa^{q-2-\theta} M_\theta. \tag{7}$$

Recall that $t_\kappa(X_{ij}) = (X_{ij} \wedge \kappa) \vee (-\kappa)$. Then for any $j = 1, \ldots, p$, as $\kappa \to \infty$, we have

$$\begin{aligned} \big|\mathbb{E}|\mathbb{E}_0[t_\kappa(X_{ij})]|^2 - \mathbb{E}|X_{ij}|^2\big| &\le \mathbb{E}|X_{ij}|^2\mathbf{1}\{|X_{ij}| \ge \kappa\} + (\mathbb{E}|X_{ij}|\mathbf{1}\{|X_{ij}| \ge \kappa\})^2 \\ &\le \kappa^{-\theta} M_\theta + \kappa^{-2(1+\theta)} M_\theta^2 \to 0, \end{aligned}$$

which implies $\min_{j\le p} \mathbb{E}|\mathbb{E}_0[t_\kappa(X_{ij})]|^2 \ge b - o(1) > 0$. Define $L_\kappa = \max_{j\le p} \mathbb{E}|\mathbb{E}_0[t_\kappa(X_{ij})]|^3$. By a similar argument as above, for any $j = 1, \ldots, p$,

$$\begin{aligned} \mathbb{E}|t_\kappa(X_{ij})|^3 &= \mathbb{E}|X_{ij}|^3\mathbf{1}\{|X_{ij}| \le \kappa\} + \kappa^3\mathbb{E}\mathbf{1}\{|X_{ij}| \ge \kappa\} \\ &\le \kappa^{1-\theta}\mathbb{E}|X_{ij}|^{2+\theta} \le \kappa^{1-\theta} M_\theta. \end{aligned}$$

So we have $L_\kappa \le 4\kappa^{1-\theta} M_\theta := \bar{L}_\kappa$. Let ϕ_κ be the quantity defined in (26) with $\bar{L}_n$ replaced by $\bar{L}_\kappa$. We are to apply Lemma 4.1 for the i.i.d. vectors $\mathbb{E}_0[t_\kappa(X_i)]$, $1 \le i \le n$, and evaluate the quantities therein. Let κ satisfy $2\kappa < \sqrt{n}/(4\phi_\kappa \log p)$. It can be easily seen that

$$\begin{aligned} M_{n,\mathbb{E}_0[t_\kappa(X)]}(\phi_\kappa) &:= \mathbb{E}\Big[\max_{j\le p} |\mathbb{E}_0[t_\kappa(X_{1j})]|^3 \mathbf{1}\Big\{\max_{j\le p} |\mathbb{E}_0[t_\kappa(X_{1j})]| > \sqrt{n}/(4\phi_\kappa \log p)\Big\}\Big] \\ &= 0. \end{aligned}$$

Let $Y_\kappa = (Y_{\kappa,1}, \ldots, Y_{\kappa,p})^\top \in \mathbb{R}^p$ be the analogue centered Gaussian vector with the same covariance matrix as $\mathbb{E}_0[t_\kappa(X_i)]$. It follows that

$$M_{n,Y_\kappa}(\phi_\kappa) := \mathbb{E}\Big[\max_{j\le p} |Y_{\kappa,j}|^3 \mathbf{1}\Big\{\max_{j\le p} |Y_{\kappa,j}| > \sqrt{n}/(4\phi_\kappa \log p)\Big\}\Big]$$

$$
\begin{aligned}
&\leq \mathbb{E}|Y_\kappa|_\infty^3 \mathbf{1}\{|Y_\kappa|_\infty > 2\kappa\} \\
&\lesssim \mathbb{P}(|Y_\kappa|_\infty > 2\kappa)\kappa^3 + \int_{2\kappa}^{\infty} \mathbb{P}(|Y_\kappa|_\infty > x) x^2 dx =: \mathrm{I} + \mathrm{II}.
\end{aligned}
$$

As has been assumed, $\mathbb{E}|Y_{\kappa,j}|^2 = \mathbb{E}|\mathbb{E}_0[t_\kappa(X_{ij})]|^2$ is upper bounded by $M_\theta^{2/(2+\theta)}$. We have $\mathrm{I} \lesssim p\kappa^3 e^{-C\kappa^2/M_\theta^{2/(2+\theta)}}$. Also, elementary manipulation implies $\mathrm{II} \lesssim p\kappa M_\theta^{2/(2+\theta)} e^{-C\kappa^2/M_\theta^{2/(2+\theta)}} + pM_\theta^{3/(2+\theta)} e^{-C\kappa^2/M_\theta^{2/(2+\theta)}}$. Hence,

$$
\begin{aligned}
M_{n,Y_\kappa}(\phi_\kappa) &\lesssim p\kappa^3 e^{-C\kappa^2/M_\theta^{2/(2+\theta)}} + p\kappa M_\theta^{2/(2+\theta)} e^{-C\kappa^2/M_\theta^{2/(2+\theta)}} + pM_\theta^{3/(2+\theta)} e^{-C\kappa^2/M_\theta^{2/(2+\theta)}} \\
&\lesssim p(nM_\theta)^{3/(2+\theta)} \exp\{-C(n^2 M_\theta^{-(2+\theta)}/\log^{4+\theta} p)^{1/(2+\theta)} \log(pn) M_\theta\} \leq n^{-1},
\end{aligned}
$$

where the second line follows in view of (5). By Lemma 4.1, we have

$$
\rho_{n,\kappa}^* := \sup_{t\in\mathbb{R}} |\mathbb{P}(\sqrt{n}|\mathbb{E}_0[\hat{\mu}_\kappa]|_\infty \leq t) - \mathbb{P}(|Y_\kappa|_\infty \leq t)| \lesssim \Big(\frac{M_\theta^2 \log^{4+3\theta} p}{n^\theta}\Big)^{1/(4+2\theta)} + \frac{1}{n}. \tag{8}
$$

With (8), we are to consider the error bound between $|\hat{\mu}_\kappa|_\infty$ and $|\mathbb{E}_0[\hat{\mu}_\kappa]|_\infty$. Since $\mathbb{E}X_i = 0$, we have

$$
\begin{aligned}
|\mathbb{E}\hat{\mu}_k|_\infty &= \max_{j\leq p} \big|\mathbb{E}(X_{ij} - \kappa)\mathbf{1}\{X_{ij} > \kappa\} + \mathbb{E}(X_{ij} + \kappa)\mathbf{1}\{X_{ij} < -\kappa\}\big| \\
&\leq \max_{j\leq p} \mathbb{E}|X_{ij}|\mathbf{1}\{|X_{ij}| > \kappa\} \leq \kappa^{-(1+\theta)} M_\theta.
\end{aligned}
$$

By (7) and Lemma 2.1 in Chernozhukov et al. (2013), for any $\delta > 0$,

$$
\begin{aligned}
\rho_{n,\kappa}^\diamond &:= \sup_{t\in\mathbb{R}} \big|\mathbb{P}(\sqrt{n}|\hat{\mu}_\kappa|_\infty \leq t) - \mathbb{P}(\sqrt{n}|\mathbb{E}_0[\hat{\mu}_\kappa]\big|_\infty \leq t)| \\
&\leq \mathbb{P}(\sqrt{n}|\hat{\mu}_\kappa - \mathbb{E}_0[\hat{\mu}_\kappa]|_\infty > \delta) + \sup_{t\in\mathbb{R}} \mathbb{P}(|\sqrt{n}|\mathbb{E}_0[\hat{\mu}_\kappa]|_\infty - t| \leq \delta) \\
&\leq \mathbb{P}(\sqrt{n}|\mathbb{E}\hat{\mu}_\kappa|_\infty > \delta) + \rho_{n,\kappa}^* + \sup_{t\in\mathbb{R}} \mathbb{P}(||Y_\kappa|_\infty - t| \leq \delta) \\
&\leq \rho_{n,\kappa}^* + 2\sqrt{n} M_\theta \kappa^{-(1+\theta)} \sqrt{\log p},
\end{aligned} \tag{9}
$$

where the last inequality follows by taking $\delta = \sqrt{n} M_\theta \kappa^{-(1+\theta)}$.
Next we compare $|Y_k|_\infty$ and $|Y|_\infty$. Observe that

$$
\begin{aligned}
\max_{j,l} |\mathrm{Cov}(Y_{\kappa,j}, Y_{\kappa,l}) - \mathrm{Cov}(Y_j, Y_l)| &= \max_{j,l} |\mathrm{Cov}(t_\kappa(X_{ij}), t_\kappa(X_{il}) - \mathrm{Cov}(X_{ij}, X_{il})| \\
&\leq C\kappa^{-\theta} M_\theta.
\end{aligned}
$$

By Lemma 3.1 in Chernozhukov et al. (2013), we obtain

$$
\rho_{n,\kappa}^\circ := \sup_{t\in\mathbb{R}} \big|\mathbb{P}(|Y_\kappa|_\infty \leq t) - \mathbb{P}(|Y|_\infty \leq t)\big| \lesssim (M_\theta/\kappa^\theta)^{1/3} (1 \vee \log(p\kappa^\theta))^{2/3}. \tag{10}
$$

Therefore, by (8), (9) and (10), we have

$$\begin{aligned}
\rho_{n,\kappa} &\le \rho^*_{n,\kappa} + \rho^\diamond_{n,\kappa} + \rho^\circ_{n,\kappa} \\
&\lesssim \rho^*_{n,\kappa} + \sqrt{n} M_\theta \kappa^{-(1+\theta)} \sqrt{\log p} + (M_\theta/\kappa^\theta)^{1/3} \log^{2/3} p \\
&= \rho^*_{n,\kappa} + \Big(\frac{M_\theta^2 \log^{4+3\theta} p}{n^\theta}\Big)^{1/(4+2\theta)} + \Big(\frac{M_\theta^2 \log^{4+3\theta} p}{n^\theta}\Big)^{1/(6+3\theta)} \\
&\le \frac{1}{n} + 3\Big(\frac{M_\theta^2 \log^{4+3\theta} p}{n^\theta}\Big)^{1/(6+3\theta)} \to 0.
\end{aligned}$$

□

Remark 1. By Propsition 2.2, the Gaussian approximation $\rho_n \to 0$ is valid for the traditional sample mean vector if each coordinate has finite q-th moment and

$$p(\log p)^{3q/2-1} = o(n^{q/2-1}). \tag{11}$$

The above condition is optimal up to a logarithmic factor. In fact, if

$$n^{q/2-1} = o(p(\log p)^{-2-q/2}), \tag{12}$$

and X_{ij} are i.i.d. symmetric random variables with $\mathbb{E}X_{ij} = 0$, $\mathbb{E}|X_{ij}|^2 = 1$ and the tail probability $\mathbb{P}(X_{ij} \ge x) = x^{-q}(\log x)^{-2}$, $x \ge x_0$, then

$$\rho_n \to 1; \tag{13}$$

see the discussion in Remark 2 of Zhang and Wu (2017). Here using the truncated sample mean, Theorem 2.1 allows p to grow exponentially with n. For example, with only finite third moment, i.e., if $M_1 = \max_{j\le p} \mathbb{E}|X_{ij}|^3 = O(1)$, p can be as large as $e^{o(n^{1/7})}$.

Proposition 2.2. *Assume (2) and there exists some constant $b > 0$ such that $Var(X_{ij}) \ge b$ for all $j = 1, \dots, p$. Then $\rho_n \to 0$ as $n \to \infty$ under*

$$p(\log p)^{3q/2-1} = o(n^{q/2-1}). \tag{14}$$

Proof. We apply Lemma 4.1. Recall the definitions of the quantities ϕ_n, $M_{n,X}(\phi_n)$ and L_n therein. Denote $\tau = \sqrt{n}/(4\phi_n \log p)$ and substitute $\phi_n = K_2(\bar{L}_n^2 \log^4 p/n)^{-1/6}$ with some $\bar{L}_n \ge L_n$. We can obtain

$$\tau = (n\bar{L}_n/\log p)^{1/3}/(4K_2) \tag{15}$$

Since $\max_{j\le q}\mathbb{E}|X_{ij}|^q<\infty$, by the Bonferroni technique and Markov's inequality,

$$M_{n,X}(\phi_n)=\mathbb{P}(|X_i|_\infty>\tau)\tau^3+3\int_\tau^\infty \mathbb{P}(|X_i|_\infty>x)x^2dx\lesssim p\tau^{3-q}. \tag{16}$$

Choose $\bar{L}_n=\max_{j\le p}\mathbb{E}|X_{1j}|^3\cdot(p^{3/q}(\log p)^{1-3/q}n^{3/q-1})^{1-\chi}(n^{1/2}(\log p)^{-7/2})^{\chi}$, where $(1-3/q)/(3/2-3/q)<\chi<1$ so that $\bar{L}_n\ge L_n$ is ensured. By (15) and (16), elementary calculation indicates that $\rho_n\to 0$ under (14). □

3 Construction of Simultaneous Confidence Intervals

Based on the Gaussian approximation result for the truncated sample mean (cf. Theorem 2.1), we are able to construct simultaneous confidence intervals (SCIs) of μ. Given a confidence level $\alpha\in(0,1)$, the $(1-\alpha)$ SCIs can be constructed by

$$\mathcal{C}_{\alpha,\kappa}=\big\{\nu=(\nu_1,\dots,\nu_p)^\top\in\mathbb{R}^p:\big|\sum_{i=1}^n t_\kappa(X_i-\nu)\big|_\infty\le\sqrt{n}c_{1-\alpha}\big\} \tag{17}$$

where $c_{1-\alpha}$ is the cutoff value determined by (19) below. Note that for every $j=1,\dots,p$, $f_j(y):=\sum_i t_\kappa(X_{ij}-y)$ is a non-increasing and continuous function of y and it is lower and upper bounded by $-n\kappa$ and $n\kappa$ respectively. Assume $n\kappa>\sqrt{n}c_{1-\alpha}$. Let l_j,u_j be the solutions to the equation $f_j(y)=\pm\sqrt{n}c_{1-\alpha}$. Then the SC for μ_j is $[l_j,u_j]$, $1\le j\le p$.

Note that $|\sum_i t_\kappa(X_i-\nu)|_\infty=0$ gives the Huber estimate.

Similarly, the null hypothesis $H_0:\mu=\mu_0$ can be rejected at level α if $|\sum_i t_\kappa(X_i-\mu_0)|_\infty\ge\sqrt{n}c_{1-\alpha}$.

Now we shall determine the cutoff value $c_{1-\alpha}$ so that the SCIs given by (17) is asymptotically valid. If $\mathrm{Cov}(X_i)=\Sigma$ is known, by Theorem 2.1, the result

$$\sup_{t\in\mathbb{R}}\Big|\mathbb{P}\Big(\frac{1}{\sqrt{n}}\Big|\sum_{i=1}^n t_\kappa(X_i-\mu)\Big|_\infty\le t\Big)-\mathbb{P}(|Y|_\infty\le t)\Big|\to 0 \tag{18}$$

implies that we can compute $c_{1-\alpha}$ by

$$c_{1-\alpha}=\inf\{t\in\mathbb{R}:\mathbb{P}(|Y|_\infty\le t)\ge 1-\alpha\}. \tag{19}$$

When Σ is unknown, one natural approach is to use a consistent estimator. Estimation of covariance matrices in high dimensions is highly nontrivial. The restriction mainly lies in the requirement of extra structural assumptions such as bandedness or sparsity.

Now we propose an innovative method to estimate the cutoff value $c_{1-\alpha}$ which can avoid the estimation of covariance matrices and is very convenient to implement. For simplicity, suppose the sample size is even denoted by $n = 2m$. Let $\pi = (\pi(1), \ldots, \pi(n))$ be a permutation of $[n] = \{1, 2, \ldots, n\}$ and define $Z_{\pi,i} = (X_{\pi(i)} - X_{\pi(m+i)})/\sqrt{2}$ for $i = 1, \ldots, m$. By such construction, $Z_{\pi,1}, \ldots, Z_{\pi,m}$ are then i.i.d. symmetric random vectors with covariance matrix Σ. By Theorem 2.1 and the fact that $\mathbb{E}t_\kappa(Z_{\pi,i}) = 0$, we have

$$\sup_{t\in\mathbb{R}} \left|\mathbb{P}(\sqrt{m}|\bar{Z}_{\pi,\kappa}|_\infty \le t) - \mathbb{P}(|Y|_\infty \le t)\right| \to 0, \tag{20}$$

where $\bar{Z}_{\pi,\kappa} = \sum_{i=1}^m Z_{\pi,i}/m$. The result (20) provides theoretical support for the validity of the procedure estimating $c_{1-\alpha}$ by the $(1-\alpha)$ quantile of $\sqrt{m}|\bar{Z}_{\pi,\kappa}|$. Let Π_n be the collection of all permutations of $[n]$ and let $\pi_1, \ldots, \pi_J$ be i.i.d. uniformly sampled from Π_n. Assume that the sampling process $(\pi_i)_{i\ge1}$ is independent of (X_i). Define

$$F_J(t) = \frac{1}{J}\sum_j^J \mathbf{1}\{\sqrt{m}|\bar{Z}_{\pi_j,\kappa}|_\infty \le t\}, \tag{21}$$

where $\bar{Z}_{\pi_j,\kappa} = \sum_{i=1}^m Z_{\pi_j,\kappa}/m$. We can obtain the empirical $(1-\alpha)$-th quantile of $F_J(t)$ by $\hat{q}_{1-\alpha} = \inf\{t \in \mathbb{R} : F_J(t) \ge 1-\alpha\}$ and estimate $c_{1-\alpha}$ by $\hat{q}_{1-\alpha}$. The simultaneous confidence interval for μ can be constructed as

$$\mathcal{C}^\dagger_{\alpha,\kappa} = \big\{\nu \in \mathbb{R}^p : \big|\sum_i t_\kappa(X_i - \nu)\big|_\infty \le \sqrt{n}\hat{q}_{1-\alpha}\big\}. \tag{22}$$

4 Appendix

Lemma 4.1 follows from Theorem 2.1 in Chernozhukov et al. (2014) for the i.i.d. case, which is useful to prove Theorem 2.1. For the i.i.d. random vectors X_i, define $L_n = \max_{j\le p}\mathbb{E}|X_{ij}|^3$ and for $\phi > 1$, define

$$M_{n,X}(\phi) = \mathbb{E}\Big[\max_{j\le p}|X_{1j}|^3\mathbf{1}\Big\{\max_{j\le p}|X_{1j}| > \sqrt{n}/(4\phi\log p)\Big\}\Big]. \tag{23}$$

Let Y be a centered Gaussian vector with $\mathrm{Cov}(Y) = \Sigma$. We can define $M_{n,Y}(\phi)$ similarly with X_{1j} replaced by Y_j in (23). Let

$$M_n(\phi) := M_{n,X}(\phi) + M_{n,Y}(\phi). \tag{24}$$

Lemma 4.1. *Suppose that there exists some constant $b > 0$ such that $\min_{j \le p} \mathbb{E} X_{ij}^2 \ge b$. Then there exist constants $K_1, K_2 > 0$ depending only on b such that for every constant $\bar{L}_n \ge L_n$,*

$$\rho_n \le K_1 \left[\Big(\frac{\bar{L}_n^2 \log^7 p}{n} \Big)^{1/6} + \frac{M_n(\phi_n)}{\bar{L}_n} \right] \tag{25}$$

with

$$\phi_n = K_2 \Big(\frac{\bar{L}_n^2 \log^4 p}{n} \Big)^{-1/6}. \tag{26}$$

References

Victor Chernozhukov, Denis Chetverikov, and Kengo Kato. Gaussian approximations and multiplier bootstrap for maxima of sums of high-dimensional random vectors. *The Annals of Statistics*, 41(6):2786–2819, 2013. 1, 4

Victor Chernozhukov, Denis Chetverikov, and Kengo Kato. Central limit theorems and bootstrap in high dimensions. *arXiv preprint arXiv:1412.3661*, 2014. 1, 7

Jianqing Fan, Weichen Wang, and Ziwei Zhu. Robust low-rank matrix recovery. *arXiv preprint arXiv:1603.08315*, 2016. 2

Danna Zhang and Wei Biao Wu. Gaussian approximation for high dimensional time series. *The Annals of Statistics*, To appear, 2017. 5